



ELSEVIER

INTERNATIONAL
JOURNAL OF SURGERYwww.theijs.com

REVIEW

Vascular smooth muscle tumors: Review of the literature

Ilias Dalainas*

492 General Military Hospital, Alexandroupolis, Greece

Available online 18 March 2007

KEYWORDS

Vascular smooth muscle tumor;
Intravascular leiomyomatosis;
Angioleiomyoma;
Vascular leiomyosarcoma;
Inferior vena cava tumor

Abstract Vascular smooth muscle tumors are very rare. They can be benign or malign. Intravascular leiomyomatosis is a benign neoplasm that extends through the veins and carries significant morbidity. Angioleiomyoma is a benign neoplasm of the extremities that carries minimal morbidity. Vascular leiomyosarcomas are malign neoplasms derived from vascular smooth cells. They are usually localized to the inferior vena cava, but can also arise from the pulmonary arteries or veins or other peripheral vessels. This study reviews literature for epidemiology, clinical presentation, diagnosis and management of patients with vascular smooth muscle tumors.

© 2007 Surgical Associates Ltd. Published by Elsevier Ltd. All rights reserved.

Introduction

Vascular smooth muscle tumors are very rare and include a variety of neoplastic lesions characterized histologically by their similarity to adult smooth muscle tissue.¹ They can be benign or malignant, with an incidence of 300 benign tumors and 1 malignant tumor per 100,000 soft tissue tumors.^{1,2} Intravenous leiomyomatosis (IVL) is a benign smooth muscle neoplasm that extends non-invasively, through the venous system, increasing however the risk of morbidity and mortality.³ Angioleiomyoma is a benign tumor arising from the vascular smooth muscle and it occurs in the extremities and the head. It is also known as angio-myoma, vascular leiomyoma, or dermal angioma.⁴ Vascular

leiomyosarcomas (LMS) are unusual malign tumors that attack arteries less often than veins. Lesions of the inferior vena cava represent the largest subgroup of these neoplasms.⁵ Other primary sites are involved far less frequently. Pulmonary vein LMS are extremely uncommon and are outnumbered in the literature by those of the pulmonary artery.⁵ This study reviews literature for the epidemiology, clinical presentation, diagnosis and management of patients with IVL, angioleiomyoma, and vascular LMS.

Intravenous leiomyomatosis

Etiology

IVL is a rare smooth muscle cell tumor that grows within the venous channels without invading them.³ About 200 cases have been reported in the literature so far.³ It was first

* Tel.: +30 694 612 4340.

E-mail address: hdlns@freemail.gr

described by Durl and Horman in 1907.⁶ The exact etiology of this neoplasm is not entirely known. Two contrasting theories have been presented, both of which have supporting evidence.¹ The first one suggests that the neoplasm arises from estrogen-induced smooth muscle cell proliferation in the venous wall of the uterine veins, while the second one suggests that the neoplasm arises from uterine leiomyomas that invade the venous system.⁷ The extension of the tumor is mostly through the uterine veins and can progress along the veins into the inferior vena cava. Further extension into the right-sided cardiac chambers will lead to intracardiac leiomyomatosis. More than 100 cases of intracardiac leiomyomatosis have been reported in the English literature until now (Table 1).³

Epidemiology

Leiomyoma shows a predilection for pre-menopausal women.^{7,8} Half of the patients have had prior pregnancies.² Since 1900, only 73 cases of cardiac leiomyomatosis have been reported¹ and 60% of the reports were within the last 15 years. However, it is believed that diagnosis is still significantly underestimated³ because it is easily missed particularly at early stages when the tumor's extension remains inside the small vessels of myometrium and cannot be detected with facility. The median age of diagnosed cases is 47 years.³

Clinical presentation

Clinical onset of these tumors usually reflects the extension of the lesions. The majority of the patients present with numerous non-specific symptoms that include vaginal

bleeding, pelvic pain, dyspnea, syncope, and congestive heart failure.^{2,7} Other symptoms include fatigue, abdominal pain, ascites, peripheral edema, and deep vein thrombosis.³ Predominant cardiac symptoms have been reported in 10% of the patients,¹ and in 50% of them uterine enlargement is observed. However, the patients may be completely asymptomatic,¹ and correct diagnosis relies on a higher index of suspicion.

IVL should be suspected when extrauterine tumor is observed in the form of wormlike or nodular plugs within pelvic veins³ during operation or unexpectedly after microscopic examination of a hysterectomy specimen.

Imaging

Echocardiography

In patients with intracardiac leiomyomatosis, transthoracic echocardiography can show the presence of a free-flowing echogenic intracardiac mass, in the right cavities or in the pulmonary arteries. Tricuspid regurgitation can also be shown.⁹ Transesophageal echocardiography usually reveals an elongated, mobile, serpent-like polypoid mass proceeding from the inferior vena cava into the right atrium and ventricle, passing through the tricuspid valve.

Computed tomography and magnetic resonance

Magnetic Resonance and 16-, 32-, 64- and 128-slide multi-detector computed tomographic angiography are the most sensitive methods to reveal IVL even in early stages without intracardiac leiomyomatosis. Heterogeneous uterine mass that can be seen unilaterally or bilaterally into the iliac veins and in the inferior vena cava is the most common finding. Masses in the subhepatic region extending down to

Table 1 Cases on intracardiac leiomyomatosis from 1900 until now

No.	Reference	Cases	Localization	Surgery
1	Lam et al. Review of cases with intracardiac extension between 1900 and 2003. <i>Int J Gynecol Cancer</i> 2003;89:175–80	68	RA 31 RV 31 PA 6	None 3 Incomplete excision 17 One-stage 19 Two-stage 29
2	Burke et al. <i>Pathology</i> 2004;36:202–3	1	PA	Death before surgery
3	Saitoh et al. <i>Gynecol Obstet Invest</i> 2004;58:168–70	1	RA	One-stage
4	Lam et al. ³	2	RA	Two-stage
5	Sasaki et al. <i>Nippon Naika Gakkai Zasshi</i> 2004;93:142–4	1	RA	One-stage
6	Jerez-Anera et al. <i>Rev Esp Anesthesiol Reanim</i> 2004;51:40–3	1	RA	One-stage
7	Sakamoto et al. <i>Jpn Thorac Cardiovasc Surg</i> 2004;52:148–51	1	RV	One-stage
8	Nishizawa et al. <i>J Am Coll Surg</i> 2004;198:842–3	1	RA	One-stage
9	Feng et al. <i>Chin Med Sci J</i> 2004;19:55	1	RA	Two-stage
10	Uchinca et al. <i>Obstet Gynecol</i> 2004;103:1068–70	1	RV	One-stage
11	Topcuoglu et al. <i>Ann Thorac Surg</i> 2004;78:330–2	1	RV	Two-stage
12	DeRubertis et al. <i>J Vasc Surg</i> 2004;40:554–8	1	RA	Two-stage
13	Bennett et al. <i>Nat Clin Pract Cardiovasc Med</i> 2005;2:369–72	1	RV	One-stage
14	Murphy et al. <i>EJVES Extra</i> 2005;9:4–6	1	RA	One-stage
15	Moorjani et al. <i>J Card Surg</i> 2005;20:382–5	1	RV	One-stage
16	Thukkani et al. <i>Ann Thorac Surg</i> 2005;79:707–9	1	RV	Two-stage
17	Ozer et al. <i>Echocardiography</i> 2005;22:514–6	1	RA	Two-stage
18	Kocica et al. ⁶	1	RV	Two-stage
19	Castelli et al. ¹⁰	1	RA	Two-stage

RA, right atrium; RV, right ventricle.

the pelvis, tumor masses within the ovarian veins or the renal veins have also been reported.³

Macroscopical, immunological, and histological findings

Excised tumors are often rubbery with a tan to whitish appearance.⁷ The largest lesion ever reported was 35 cm in length.⁷ IVL are often free-floating and elongated. Grossly, the lesion is characterized by coiled masses within the myometrium and serpentine extensions into the uterine veins.^{1,2}

In immunological studies the tumor cells are diffusely and strongly positive on smooth muscle markers, including desmin and smooth muscle actin,³ confirming their smooth muscle nature. They show bizarre nuclear morphology with hyperchromatic multilobulated nuclei. The mitotic activity is low, with mitotic index of less than 1 per 50 high-power fields. The cells range from highly cellular spindle cell masses to cellular ones marked by fibrosis, hydropic change, and perivascular hyalinization.^{1,2} Tumors positive for CD34 are often non-thrombogenic due to the presence of an endothelial surface layer.^{1,7}

Several reports included immunohistochemically positive reactions of the intravascular parts of the tumor⁹ with antibodies against estrogen and progesterone receptors, and it was suggested that a high concentration of serum estradiol and a high level of tissue estrogen receptor are related to the IVL and potentially influence the growth of the smooth muscle cells.^{9,10} The latter theory could be potentially confirmed because the highest incidence rate of recurrence is found in patients with normal ovarian activity. It was also suggested that occasionally hormonal therapy should be considered in cases of unresectable residual tumors.⁹ The detection of cytoplasmic estradiol and progesterone receptors in both the intravascular tumor and leiomyosarcoma would allow consideration of the use of an antiestrogen such as Tamoxifen to control the progress of the disease, although there is not any report of efficacy in the literature.⁹

Treatment

Surgical resection is the treatment of choice.^{1,2,7,10} Attempts must be made to remove the entire neoplasm, which usually involves hysterectomy.¹ Incomplete excision can result in recurrence of the neoplasm. Recurrence rate of 30% from 7 months to 17 years follow-up has been reported.^{2,7} In a study by Butany et al.¹ there was a 12.5% of recurrence after complete excision in 8.6 months' follow-up. When treating IVL, even if risk of pulmonary embolism is low, a temporary inferior vena cava filter can prevent from potential embolism during surgical maneuvers. When both abdominal and thoracic approach is needed, a two-stage operation can be considered. The slow growth of the tumor allows a safe interval between two major surgical procedures. The advantages of two-stage strategy are the shorter operative times, despite the risks of a second general anesthesia, and the less risk of bleeding, as systemic heparinization mandatory for one-stage cardiopulmonary bypass, is actually avoided. In the thoracic approach the tumor should be

resected below the level of the renal veins to facilitate easier vascular control in the second-stage laparotomy,³ because higher vascular control requires mobilization of the liver via a right thoracoabdominal incision, with its associated morbidities. In addition, clamping of the cava above the hepatic vein congruence will significantly reduce the preload, necessitating an axillofemoral venous bypass.³

It has been suggested that recurrence may show the same pattern even after hysterectomy and bilateral adnexectomy. This shows that the tumor growth is independent of the presence of the uterus and although histologically benign, might be considered clinically malignant.¹⁰ When recurrence is seen, reintervention is actually universally recommended to achieve long-term disease-free survival.¹⁰

Angioleiomyoma

Etiology/epidemiology

Minor trauma, venous stasis and hormonal changes, especially estrogenic, have been proposed as etiological features. The presence of chronic inflammatory infiltrates in some lesions supports the venous stasis theory.⁴ Another theory suggests that the lesions could be hamartomatous as has been demonstrated by the presence of mature fat cells.⁴

They account for 5% of all benign neoplasms of soft tissues and their prevalence in eastern South Africa is 10 times that of the Caucasian population living in temperate climates.⁴

Histology

In 1973 Morimoto¹¹ studied 241 cases of angioleiomyoma and classified them into three histological types:

1. Solid: The most common type which has closely compacted smooth muscle and many small, split-like vascular channels. This type of tumors is three times more common in females than in males.
2. Venous: Thick, easily identifiable muscular walls distinguish this type. These tumors occur more commonly in males.
3. Cavernous: The vascular channels are dilated with less smooth muscles. This is the most uncommon type. These tumors occur 4 times more common in males than in females.

Morimoto¹¹ also grouped these tumors into two groups: the larger group of extremity tumors, where they are mainly of the solid type, often painful, and the smaller group of head tumors, where they are usually of the venous type and painless.

Clinical presentation

The typical finding is a solitary, small, slow-growing, firm, mobile, subcutaneous nodule. Most of them are less than 2 cm in size.⁴ Pain is the most striking clinical feature of angioleiomyoma. It is often paroxysmal and is provoked by exposure to cold and wind.⁴

Imaging

On ultrasound, it shows well defined margins and a homogeneous structure, suggesting benign nature. Color-duplex shows high resistance in intratumor arteries, suggesting the presence of muscular arteries.⁴ Magnetic resonance shows the tumors hypervascular and isointense to skeletal muscle. For an extremity mass with mixed areas of hyper and hypointensity to skeletal muscle of T2-weighted images and hypointense rim, a diagnosis of muscular leiomyoma should be considered.⁴

Treatment

Simple excision for biopsy is often curative.⁴ Recurrences have occasionally been reported.⁴

Leiomyosarcoma

Etiology/epidemiology

Unlike IVL, LMS are malignant neoplasms derived from vascular smooth cells. They represent about 5–7% of all soft tissue sarcomas.^{12–15} They most often affect the gastrointestinal system and the uterus, less frequently the respiratory apparatus, and rarely the limbs.¹⁶ The adult population is most susceptible, particularly the women.^{12–15} A theory for the explanation of female predominance is that growth and proliferation of the smooth muscle is likely to be influenced by pregnancy and estrogenic stimulation. LMS can be distinguished as intraperitoneal and retroperitoneal, cutaneous, subcutaneous and vascular neoplasia.¹⁶

Vascular LMS represent only 2% of all these neoplasias and about 200 cases have been reported in the literature.^{16–18} Veins are involved five times more often than arteries and particularly the vena cava, which alone represents 50% of all localizations, followed by the big systemic veins and the great saphenous.¹⁹ Arterial LMS are equally distributed between pulmonary artery (the most frequent localization, approximately 50% of the cases¹⁶) and other systemic arteries. Review of the published literature identified 19 reported cases of peripheral (non pulmonary/non aortic) arteries LMS.^{17,19–36} They refer to 9 women and 10 men. These 19 patients ranged from 33 to 79 years old with an average age of 58.6 years. Since the initial report by Perl in 1871, there have been 218 cases of LMS of the inferior vena cava reported,^{2,37,38} and over the last 18 years 70.4% of reported LMS occurred in the inferior vena cava.¹ The male to female ratio is 3:7 with a mean age of 56.9 ± 7.9 years.¹ LMS have been reported to arise in the pulmonary arteries and veins with a ratio of 20:1.⁵ Pulmonary vessel LMS has a male to female ratio of 1:2.¹ Four cases of LMS of the portal and large mesenteric veins have been reported³⁹ in the literature until now.

Clinical presentation

Abdominal pain is the most common symptom for neoplasms occurring in any region of the body and is referred from 45.7% of the patients.¹ Deep vein thrombosis or venous distension are observed in 12% and 20% of the patients

respectively. Inferior limb edema is also a common symptom.⁴⁰ When the localization is a pulmonary vessel, symptoms like palpitation, dizziness, syncopal attacks, dyspnea and eventual right-heart failure can be present.^{2,38} In peripheral artery localization the main symptoms are pain and vascular insufficiency. However, in approximately one third of the patients the diagnosis is incidental as the patients are asymptomatic. Therefore, the diagnosis is often delayed, despite the tumor's large size, and it remains asymptomatic for a long time.

Diagnosis

Common laboratory tests reveal albuminuria when the hepatic or renal veins are involved. Polycythemia and microangiopathic anemia can also be observed.⁴¹ Gastrointestinal and intravenous contrast studies demonstrate an extrinsic mass effect when the localization is the inferior vena cava. In ultrasonography of the abdomen the tumor appears as a mass with homogeneous or heterogeneous sound patterns.^{42–53} Doppler ultrasound gives information about patency of the portal or systemic venous system.^{43,44,50}

Computed tomography and magnetic resonance

Computed tomography and magnetic resonance are the most sensitive examinations and confirm the presence of the tumor, its pattern or growth, relationship to the surrounding structures, and the presence of caval obstruction. On computed tomography the tumor appears as low density, solid, or heterogeneous mass compressing or silhouetting, involving or arising from the inferior vena cava.^{41,43,54–61} Magnetic resonance gives similar information but the sagittal section is more informative as to the extent of the tumor and tumor thrombosis.^{41,45,50,52,53,60–62}

Angiography

On angiography, LMS exhibit variable degrees of vasculability.^{43,55,57,62–64} Hypervascular tumors derive their blood supply from the hepatic artery or its branches, adrenal, lumbar, gastric, pancreatoduodenal, or gastroduodenal artery. Other findings can be distortion, stretching, or compression to adjacent structures. Angiography cannot differentiate caval LMS from retroperitoneal LMS. In arterial localization of LMS, the arteriogram may confuse it with pseudoaneurysms.³⁶

Histology

LMS of the inferior vena cava exhibits a typical pattern of interlacing sweeping bundles of spindle-shaped cells with elongated, blunt-ended nuclei with a tendency to palisade.⁴¹ Mitotic figures can vary within the same tumor. Mononuclear giant cells can be present at a variable degree, and areas of necrosis are commonly seen. Masson's trichrome and Van Gieson's stain indicate the presence of collagen tissue. Immunohistochemical and ultrastructural studies reflect features found in soft tissue sarcoma of other sites. The tumor cells display desmin, vimentin, and smooth cell actin, but not S-100 protein.^{41,58,65,66} A high desmin immunoreactivity rate exceeding that seen in other soft tissue LMS is also noted.⁴¹ Myofibrils, intracytoplasmic vesicles, and occasional tight junctions between tumor cells are described.^{41,42,66,67}

Biological behavior

LMS of the inferior vena cava are considered to be slowly growing tumors. In a review of 89 resected and 43 non-resected LMS of the inferior vena cava,⁴¹ three cases of rather rapid growth rate were identified.^{62,64,68} In these cases the tumor was not present on previous laparotomy performed 6–24 months for unrelated disorders.

The tumor metastasize late in the course of the disease and spreads mainly systemically and most commonly to the lungs and the liver, but no organ is exempt. Lymph node metastasis occurs less frequently.⁴¹

Management

Surgical

Complete surgical resection is the mainstay of treatment for vascular LMS.⁶⁹ Whenever possible, surgical resection should be used to completely remove the primary tumor intact and en bloc with any directly involved adjacent structures that can be sacrificed. Ideally, a margin of uninvolved normal tissue would be resected with the tumor, given the high incidence of local recurrence when lesser resections are performed.³⁶ The necessary disruption of the vascular system will usually require vascular reconstruction after resection. The location of the primary tumor and its extent will dictate the type and complexity of the reconstruction required to reestablish arterial flow or caval flow. Tumor thrombectomy alone, even if not curative, can prevent death from liver failure and right heart outflow tract obstruction in some cases.⁴¹

Radiation

Adjuvant radiation therapy, either pre- or postoperatively, decreases the rates of local recurrence of LMS.³⁶ Some authors^{69,70} suggested that high-dose-rate intraoperative radiation improves local control; however, the utility of radiation therapy in this setting should be ultimately be determined by prospective randomized trials.

Chemotherapy

Effective systemic chemotherapy for the treatment of LMS of soft tissues is at least controversial.³⁶ Toxicity is significant, and systemic chemotherapy would only be considered for large high-grade tumors, recurrent tumors, and metastatic disease. Appropriate follow-up for these tumors should monitor for focal recurrence (with CT or MRI every 3–4 months³⁶) and for pulmonary metastasis (with CT of chest every 4–6 months³⁶) during the first 5 years postoperatively.

Prognosis

Prognosis following complete resection varies according to the histologic features of the primary tumor. In a review of inferior vena cava LMS,⁴¹ in Grade I (with minimal, few, or less than five mitotic figures per high-power field), survival with no evidence of disease for 3 months to 13 years was noticed in 64.3% of the patients. In grade II (moderately differentiated tumor), survival with no evidence of the disease for 9 months to some years was observed in 60% of the patients, and local recurrence developed 20%. In grade

III (poorly differentiated tumor, with frequent or more than 10 mitotic figures per high-power field), survival with no evidence of the disease for 2–15 months was seen in 57% of the patients, whereas systemic spread occurred within 3 months to 2 years in 43%. However, other authors⁶² did not find a correlation between mitotic activity and clinical outcome.

Conclusions

Vascular smooth muscle neoplasms that extend through blood vessels are extremely rare, difficult to be diagnosed early, and complex to treat.

IVL is a benign smooth muscle cell tumor that grows within the veins. It is a complex disease that carries significant morbidity, and multidisciplinary surgical approach is the key for an accurate tumor excision.

Angioleiomyoma is a benign tumor which commonly presents between the third and fifth decade of life. It is painful in over half of the cases. It should be considered in the differential diagnosis of painful nodular lesions of the extremity. It causes minimal morbidity and excision is usually curative.

Vascular LMS are potentially aggressive neoplasms. The management of these tumors is today oriented toward preoperative staging and complete surgical resection with vascular reconstruction. However, prognosis depends also on the size and the grade of the tumor. The only therapeutic modality associated with prolonged survival is complete surgical resection with clean margins, followed by adjuvant radiation therapy.

Conflicts of interest

None.

Funding

None.

Ethical approval

Not requested.

References

1. Butany J, Singh G, Henry J, Leong SW, Nair V, Thangaroopan M, et al. Vascular smooth muscle tumors: 13 cases and a review of the literature. *Int J Angiol* 2006;**15**:43–50.
2. Enzinger F, Weiss S. In: Gay S, editor. *Soft tissue tumors*. St. Louis: Mosby Yearbook; 2001. p. 1120.
3. Lam PM, Lo KW, Yu MY, Wong WS, Lau JY, Arifi AA, et al. Intravenous leiomyomatosis: two cases with different routes of tumor extension. *J Vasc Surg* 2004;**39**:465–9.
4. Ramesh P, Annapureddy SR, Khan F, Sutaria PD. Angioleiomyoma: a clinical, pathological and radiological review. *Int J Clin Pract* 2004;**58**:587–91.
5. Oliai BR, Tazelaar HD, Lloyd RV, Doria MI, Trastek VF. Leiomyosarcoma of the pulmonary veins. *Am J Surg Pathol* 1999;**23**:1082–8.
6. Kocica MJ, Vranes MR, Kostic D, Kovacevic-Kostic N, Lackovic V, Bozic-Mihajlovic V, et al. Intravenous leiomyomatosis with extension to the heart: Rare or underestimated? *J Thorac Cardiovasc Surg* 2005;**130**:1724–6.

7. Ling FT, David TE, Merchant N, Yu E, Butany JW. Intracardiac extension of intravenous leiomyomatosis in a pregnant woman: a case report and review of the literature. *Can J Cardiol* 2000; **16**:73–9.
8. Trojani M, Contesso G, Coindre JM, Rouesse J, Bui NB, de Mascarel A, et al. Soft-tissue sarcomas of adults; study of pathological prognostic variables and definition of a histopathological grading system. *Int J Cancer* 1984; **33**:37–42.
9. Baca Lopez FM, Martinez-Enriquez A, Castrejon-Aivar FJ, Ruanova-Leon D, Yanez-Gutierrez L. Echocardiographic study of an intravenous leiomyoma: case report and review of the literature. *Echocardiography* 2003; **20**:723–5.
10. Castelli P, Caronno R, Piffaretti G, Tozzi M. Intravenous leiomyomatosis with right heart extension: successful two-stage surgical removal. *Ann Vasc Surg* 2006; **20**:405–7.
11. Morimoto N. Angioleiomyoma. A clinicopathologic study. *Med J Kagoshima Univ* 1973; **24**:663–83.
12. Kokawa K, Yamoto M, Yata C, Mabuchi Y, Umesaki N. Postmenopausal intravenous leiomyomatosis with high levels of estradiol and estrogen receptor. *Obstet Gynecol* 2002; **100**:1124–6.
13. Enzinger F, Weiss S. *Soft tissue tumors*. St Louis: Mosby; 1982. p. 311–2.
14. Fisher MG, Gelb AM, Nussbaum M, Havenson S, Ghali V. Primary smooth muscle tumors of venous origin. *Ann Surg* 1982; **196**:720–4.
15. Stout A, Lattes R. *Tumors of the soft tissues. Atlas of tumor pathology. Series 2, fascicle I*. Washington, DC: Armed Forces Institute of Pathology; 1967. p. 127–33.
16. Caldarelli GF, Barellini L, Faviana P, Guerra M. Leiomyosarcoma of the popliteal artery. Case report and review of the literature. *J Vasc Surg* 2003; **37**:206–9.
17. Leeson MC, Malaee M, Makley JT. Leiomyosarcoma of the popliteal artery: a report of two cases. *Clin Orthop* 1990; **253**:225–30.
18. Rhode HT, Riesener KP, Buttner R, Schumpelick V. Leiomyosarcoma of the splenic artery. *Chirurg* 2001; **72**:844–6.
19. Stringer BD. Leiomyosarcoma of artery and vein. *Am J Surg* 1977; **134**:90–4.
20. Kevorkian J, Cento DP. Leiomyosarcoma of large arteries and veins. *Surgery* 1973; **73**:390–400.
21. Gutman H, Haddad M, Zelikovski A, Mor C, Reiss R. Primary leiomyosarcoma of the right common iliac artery—a rare finding and cause of occlusive vascular disorder. *J Surg Oncol* 1986; **32**:193–5.
22. Hopkins GB. Leriche syndrome associated with leiomyosarcoma of the right common iliac artery. *JAMA* 1968; **18**:1789–90.
23. Birkenstock WE, Lipper S. Leiomyosarcoma of the right common iliac artery: a case report. *Br J Surg* 1976; **63**:81–2.
24. Delin A, Johansson G, Silfversward C. Vascular tumors in occlusive disease of the iliac-femoral vessels. *Eur J Vasc Surg* 1990; **4**:539–42.
25. Wright EP, Glick AD, Virmani R, Page DL. Aortic intimal sarcoma with embolic metastases. *Am J Surg Pathol* 1985; **9**:890–7.
26. Varela-Duran J, Oliva H, Rosai J. Vascular leiomyosarcoma: the malignant counterpart of vascular leiomyoma. *Cancer* 1979; **44**:1684–91.
27. Mikami Y, Manabe T, Lie JT, Sakurai T, Endo K. Intramural sarcoma of the carotid artery with adventitial inflammation and fibrosis resembling “inflammatory aneurysm”. *Pathol Int* 1997; **47**:569–74.
28. vanGulik TM, Taat CW, Regroort M, Knomhout JG, Thunnissen FB. Vascular leiomyosarcoma—a report of two cases. *Neth J Surg* 1987; **39**:72–3.
29. Giangola G, Migaly J, Crawford B, Moskowitz P, Sebenick M. Leiomyosarcoma of the subclavian artery. *J Vasc Surg* 1995; **22**:496–500.
30. Meulman NB, Graham JC. Leiomyosarcoma of the superficial femoral artery presenting as a ruptured aneurysm. *Aust N Z J Surg* 1991; **61**:853–5.
31. Basu SK, Scott TD, Wilmshurst CC, MacEachern AG, Clyne CA. Leiomyosarcomata of the popliteal vessels: rare primary tumors. *Eur J Vasc Surg* 1988; **2**:423–5.
32. Briggs PJ, Pooley J, Malcolm AJ, Chamberlain J. Popliteal artery leiomyosarcoma: a case report and review of the literature. *Ann Vasc Surg* 1990; **4**:365–9.
33. Lo KWK, Lau TK. Intracardiac leiomyomatosis: case report and literature review. *Arch Gynecol Obstet* 2001; **264**:209–10.
34. Nam MS, Jeon MJ, Kim YT, Kim JW, Park KH, Hong YS. Pelvic leiomyomatosis with intracaval and intracardiac extension: a case report and review of the literature. *Gynecol Oncol* 2003; **89**:175–80.
35. Armstrong PJ, Franklin DP. Pararenal vena cava leiomyosarcoma versus leiomyomatosis: difficult diagnosis. *J Vasc Surg* 2002; **36**:1256–9.
36. Blansfield JA, Chung H, Sullivan TR, Pezzi CM. Leiomyosarcoma of the major peripheral arteries: case report and review of the literature. *Ann Vasc Surg* 2003; **17**:565–70.
37. Nikaido T, Endo Y, Nimura S, Ishikura H, Ushigome S. Dumbbell-shaped leiomyosarcoma of the inferior vena cava with foci of rhabdoid changes and osteoclast-type giant cells. *Pathol Int* 2004; **54**:256–60.
38. Gehr NR, Lund O, Alstrup P, Nielsen JS, Villadsen AB, Bartholdy NJ. Recurrence of uterine intravenous leiomyomatosis with intracardiac extension. Diagnostic considerations and surgical removal. *Scand Cardiovasc J* 1999; **33**:312–4.
39. Celdrán A, Frieyro O, del Rio A, Franco A, Bosch O, Sarasa JL. Leiomyosarcoma of the portal venous system: A case report and review of the literature. *Surgery* 2004; **134**:455–6.
40. Spencer JM, Amonette RA. Tumors with smooth muscle differentiation. *Dermatol Surg* 1996; **22**:761–8.
41. Kulaylat MN, Karakousis CP, Doerr RJ, Karamanoukian HL, O'Brien J, Peer R. Leiomyosarcoma of the inferior vena cava: a clinicopathologic review and report of three cases. *J Surg Oncol* 1997; **65**:205–17.
42. Patel JK, Englander LS. Leiomyosarcoma of the inferior vena cava. *J Surg Oncol* 1982; **21**:238–40.
43. Thompson MM, Graham TR, Bolia AA, Firmin RK, Bell PR. Intrahepatic leiomyosarcoma of the inferior vena cava with extension into the right atrium. *Eur J Vasc Surg* 1993; **7**:204–7.
44. O'Malley KJ, Stuart RC, McEntee GP. Combined resection of the inferior vena cava and extended right hepatectomy for leiomyosarcoma of the retrohepatic cava. *Br J Surg* 1994; **81**:845–6.
45. Kracht M, Becquemin JP, Anglade MC, Mathieu D, Hillion ML, Teboul JL. Acute Budd-Chiari Syndrome secondary to leiomyosarcoma of the inferior vena cava. *Ann Vasc Surg* 1989; **3**:268–72.
46. Yanaga K, Okadome K, Ito H, et al. Graft replacement of pararenal inferior vena cava for leiomyosarcoma with the use of venous bypass. *Surgery* 1993; **113**:109–12.
47. Chauhan MA, Smith PL, Ferris EJ, Murphy K, Westbrook K, Slayden JE. Leiomyosarcoma of the inferior vena cava: Angiography and computed tomography findings: Report of two cases and review of imaging criteria. *Cardiovasc Intervent Radiol* 1981; **4**:209–14.
48. Bruyninckx CMA, Derksen OS. Leiomyosarcoma of the inferior vena cava: Case report and review of the literature. *J Vasc Surg* 1986; **3**:652–6.
49. Kretz JG, Matysiak L, Matter D, Wenger JJ, Tonqio J, Immler M, et al. Sarcome leiomyoblastique de la veine cave inferieure: A propos d'un cas recent. *Ann Chir* 1984; **38**:309–11.
50. Fabre JM, Domergue J, Fagot H, et al. Leiomyosarcoma of the inferior vena cava presenting as Budd-Chiari syndrome. Vena cava replacement under veno-venous bypass and liver hypothermic perfusion. *Eur J Surg Oncol* 1995; **21**:86–7.
51. Nyman U, Jonsson K. Angiography in leiomyosarcoma of the inferior vena cava: Report of two cases. *Br J Radiol* 1979; **52**:273–5.

52. Griffin AS, Sterchi JM. Primary leiomyosarcoma of the inferior vena cava: A case report and review of the literature. *J Surg Oncol* 1987;**34**:53–60.
53. Kasano Y, Tanimura H, Kumada K, Taniguchi K, Takifuji K, Ohnishi H, et al. Resectable leiomyosarcoma of inferior vena cava extended into the right atrium with the use of cardiopulmonary bypass and graft replacement. *Surgery* 1995;**117**:473–5.
54. Fischer GM, Gelb AM, Nussbaum M, et al. Primary smooth muscle tumors of venous origin. *Ann Surg* 1982;**196**:720–4.
55. Demers ML, Curley SA, Romsdahl MM. Inferior vena cava leiomyosarcoma. *J Surg Oncol* 1992;**51**:89–93.
56. Mingoli A, Feldaus RJ, Cavallaro A, Stipa S. Leiomyosarcoma of the inferior vena cava: Analysis and search of the world literature on 141 patients and report of three new cases. *J Vasc Surg* 1991;**14**:688–99.
57. Young R, Friedman AC, Hartman DS. Computed tomography of leiomyosarcoma of the inferior vena cava. *Radiology* 1982;**145**:99–103.
58. Satoh M, Katoh J, Onodera S. Leiomyosarcoma of the inferior vena cava causing Budd-Chiari syndrome: A case report. *Angiology* 1993;**44**:673–6.
59. Case Records of the Massachusetts General Hospital: Weekly clinicopathological exercises. *N Engl J Med* 1981;**304**:162–8.
60. Monig SP, Gawenda M, Erasmi H, Zieren J, Pichlmaier H. Diagnosis, treatment and prognosis of Leiomyosarcoma of the inferior vena cava. *Eur J Surg* 1996;**4**:231–5.
61. Rosenthal JT, Colonna JO, Drinkwater DC. Leiomyosarcoma of the inferior vena cava with atrial extension: Long-term survival following resection and caval replacement with circulatory arrest. *Urology* 1995;**46**:876–8.
62. Dzsinič C, Gloviczki P, van Heerden JA, et al. Primary venous leiomyosarcoma: A rare but lethal tumor. *J Vasc Surg* 1992;**15**:595–603.
63. Tegtmeyer CJ, Buschi A. The angiographic diagnosis of leiomyosarcoma of the inferior vena cava. *Radiology* 1977;**122**:683–5.
64. Wray Jr RC, Dawkins H. Primary smooth muscle tumor of the inferior vena cava. *Ann Surg* 1971;**174**:1009–19.
65. Burke AP, Virmani R. Sarcomas of the great vessels: A clinicopathologic study. *Cancer* 1993;**71**:1761–73.
66. Taylor RW, Sylwestrowicz T, Kossakowska AE, Urbanski SJ, Minuk GY. Leiomyosarcoma of the inferior vena cava presenting as Budd-Chiari syndrome. *Liver* 1987;**7**:201–5.
67. Nickels J, Kalima TV. Leiomyosarcoma of the inferior vena cava. *VASA* 1979;**8**:333–6.
68. Vreede AA, Vaziri D, Cornwall CC. Leiomyosarcoma of the inferior vena cava. *Vasc Surg* 1982;**16**:143–7.
69. Hollenbeck ST, Grobmyer SR, Craig Kent K. Surgical Treatment and outcomes of patients with primary inferior vena cava leiomyosarcoma. *J Am Coll Surg* 2003;**197**:575–9.
70. Alektiar KM, Hu K, Anderson L, Brennan MF, Harrison LB. High-dose-rate intraoperative radiation therapy (HDR-IORT) for retroperitoneal sarcomas. *Int J Radiat Oncol Biol Phys* 2000;**47**:157–63.